REFERENCES

Abdullahi, H. A.; Azionu, B. C. and Ajayi, O. (2007): Checklist of zooplankton in culture tanks at NIFFRI Green House, New Bussa. Proceedings of the 22nd Annual Conference of Fisheries Society of Nigeria (FISON), Kebbi 12th-16th November, 2007. Vol. 1:284-290.

abundance and hydrological condition in laguna, Jogyunda, Puerto rico, 2(3): 625 – 631.

Adeyemi, S. O.; Adikwu, L. A.; Akombu, P. M. and Iyela, J. T. (2009): Survey of zooplankton and macroinvertebrates of Gbedikere Lake Bassa, Kogi State, Nigeria. *International Journal of Lakes and River,* **2**(1): 37–44.

Aguigwo, J. N. (1998): Studies on the physico-chemical parameters and plankton productivity of a productive stream. *Journal of Aquatic Science,* **13**: 9–13.

Akin–Oriola, F. A. (2003): Zooplankton associations and environmental factors in Ogunpa and Ona Rivers, Nigeria. *Rev. Biol. Trop*., **51** (2): 391–398.

Ali, S.S., 1999, Fresh Water Fisheries Biology Isted. Naseem Book, Depot. Hyderabad.

Allan, J. D. (2001): Stream Ecology: structure and function of running waters. School of Natural Resources and Environment, University of Michigan, USA. Kluwen Academic Publishers. 175pp.

American Public Health Association (APHA), American Water Work Association (AWWA) and Water Pollution Control Federation (WPCF) (2005): Standard methods for the examination of water and waste water. 21st ed. APHA, Washington D. C., 1195pp.

Anand, N. *Indian freshwater microalgae*, Bishen Singh MahendrapalSinghPublishers, 1998.

Anand, N., 1998. Freshwater Micro alagae. Bishen Singh Mahendra Pal Singh, Dehradun, India. 94.

andPadamsar lakes. Ph. D. Thesis Submittedto J. N. V. University, Jodhpur Rajasthan.

Annalakshmi .G and Amsathi . A. (2012). Studies on the hydrobiology of river cauvery and its tributaries arasalarfromkumbakonam region (tamilnadu, india) with reference to zooplankton.International journal of applied biology andPharmaceutical Technology. Volume: 3(1).

APHA (1992). *Standard Methods for the Examination of Water and Wastewater* (18th edition). American Public Health Association, Washington D.C.

APHA, AWWA and WEF. *Standard methods for the examination of waterand wastewater*. 21st edition. Published jointly by the American Public HealthAssociation, American Water Works Association and Water EnvironmentFederation, 2005. Smith, GM. *The Freshwater algae of the United States*, McGraw Hill, NewYork, USA, 1950.

Baruah, A., A.K. Sinha & U.P. Sharma (1993). Plankton variability of a tropical wetland, Kawar (Begusarai), Bihar. *Journal of Freshwater Biology* 5: 27–32.

Benarjee G, K Srikanth, G Ramu, K NarasimhaRamulu and B Ravinder (2008). "The Climatic Influence onZooplanktonic Population in Historical lake of Kakatiya Dynasty", Proc. of 8th Indian Fisheries Forum, pp. 22-26,Kolkata.

Bhagat, V.B. and C.B. Meshram (2007). Zooplankton dynamics of Ambadi dam, near Akot, dist. Akola, Maharashtra.Journal of Aquatic Biology Volume: 22(1):19-20.
BhoraChandan and Aravind Kumar (2004). Plankton diversity in the wetlands of Jharkand.A.P.H publishing Corp .NewDelhi: 91-123.
Boxshall, G. A. and E. E. Strong, (2006). An extraordinary shift in life habit in a genus of cyclopid copepods from LakeTanganyika. Zoological Journal of the Linnean Society 146: 275–285.
Boxshall, G. A. and T. D. Evstigneeva, (1994). The evolution of species flocks of copepods in Lake Baikal: apreliminary analysis. In: Martens, K., B. Goddeeris& G. Coulter (eds), Speciation in Ancient Lakes. Archivfu¨rHydrobiologie, Ergebn Limnology 44: 235–245.

Boyd, C.E., 1981. Water quality in warm water fish ponds craft master printers. Inc. Opelika, Alabama.
Coulter, G., (1991). Lake Tanganyika and its Life. Oxford University Press, Oxford.

Cox, EJ. *Identification of freshwater diatom live material*, Chapman and Hall,London, 1996.
Davies, O. A.; Abowei, J. F. N. and Otene, B. B. (2009): Seasonal abundance and distribution of plankton of Minichinda stream, Niger Delta, Nigeria. *American Journal of Scientific Research,* **2**: 20-30.
Davies, O.A., Tawari C.C. and Abowei J.F.N., ZooplanktonofElechi Creek, Niger Delta Nigeria, *Environ. Ecol*.,26(4c), 2441-2346 (2008)7. Neves I.F., Recha O., Roche K.F. and
De Graaf G.J., Marttin F. (2003). Mechanisms behind changes in fish biodiversity in the floodplains of Bangladesh. –
Desikachary, TV. *Cyanophyta*, Indian Council of Agricultural Research, NewDelhi, 1959.

Dhanapathi, M.V., (2000). Taxonomic notes on the rotifers from India (from 1889-2000). Indian Association of AquaticBiologists (IAAB), Hyderabad.
Dhanapathi, M.V.S.S.S. 2000. Taxonomic noteson the *Rotifera*, Indian Association ofAquatic Biologist, Hyderabad, Vit, 178Pp.
Dhembare, A.J. 2011. Diversity and its indices in zooplankton with physicochemical properties of Mula Dam WaterAhmednagar, Maharashtra India. *Eur. J.Exp. Biol.,* 1(4): 98 103.Edmondson, W.T. 1959. Fresh water biology,
Durand, J. R. et Leveque, C. (1980): Flora et fauneaquatiques de L´Afrique. Sahelo-SoudanienneTomme 1 Paris. 189pp.
Dutta S.P.S,H.Kaur and I.D.Sharma (2001). Hydrobiological studies on river Basantar samba (Jammu and Kashmir)J.Aqua.bio.16,41-44
Dwivedi, S., Misra, P.K., Tripathi, R.D., Rai, U.N., Dwived, C.P., Baghal, V.S., Suseela, M.R., and Srivastava, M.N., 2005. Systematic and ecological studies on chlorophyceae of North India and their relationship with water quality. *J. Environ. Biol.* 26: 495–505.
Edmendson, W.J., 1959. Fresh water biology, 2nd Edn., John Wiley and Sons, New York, 1248.

Edmondson, W. T. (1992): Fresh WaterBiology. IInd Ed. *John Wiley and Sons*. Inc. NewYork., 1248pp.
Edward AndHipple (Eds), 2nd edn. John Willy & Sons Inc., Newyork, Pp. 95189.Gaike, P., Mohite, M., Ningule, A., Kambale,A., Shejule, K. 2012. Seasonalzooplankton variation. In: Nagatas Dam near ParturDistJalna, India. *J. Biosci.Disc.*, 3(3): 362 365.

Ekpo I. E., Chude, L. A., Onuoha, G. C. and Udoh, J. P. (2012c): Studies on the physico-chemical characteristics and nutrients of a tropical rainforest river in southeast Nigeria. *AACL Bioflux*, **5**(3):142-162.
Ekpo, I. E. (2012): Diversity and condition factor of fish species of Ikpa River at Nwaniba in Niger Delta, Nigeria. *FUTA Journal of Research in Sciences,* **1**(1): 36-47.
Ekpo, I. E. (2013): Women's participation in lower Ikpa River fisheries of AkwaIbom State Nigeria: A case-study of Ifiayong. *Journal of Fisheries and Aquatic Science,* **8**(1): 268 - 278.

Ekpo, I. E. Chude, L. A. and Onuoha, G. C (2012a): The use of multivariate analysis for characterisation and classification of Ikpa River, Nigeria. *Elixir Agriculture,* **49**: 9799-9807.
Ekpo, I. E., Onuoha, G. C. and Chude, L. A. (2011): Diversity and variability of aquatic macrophytes in Ikpa River, IkotEbom (AkwaIbom State) Nigeria. *Nigerian Journal of Agriculture, Food and Environment,* **7**(2): 1-9.
Ekpo, I. E., Udo, M. T. and Usip, P. L. **(**2012b): Seasonality and size variation of fish species in Nwaniba, Ikpa River Southeast Nigeria. *Elixir Agriculture,* **51**: 11043 – 11050.
Eshwarlad, S. and Angadi, S.B. (2003). Physico-chemical parameters of two fresh water bodies of Gulbarga India withspecial reference to phytoplankton. *Poll. Res*., 22(3): 411-422.
Eswari,Y.N.K., and Ramani Bai, 2002. Distribution and abundance of phytoplankton is estuarine waters of Chennai, South East coast of India. *J. Mar. Biol.* Ass. India. 44: 205 – 211.

Fritsch, F.E., 1961. The structure and reproduction of Algae. Vol. 1. Cambridge at University Press. Cambridge.
Gaiwad, S.R., Tarot, S.R., and Chavan, T.P., 2004. Diversity of plankton and Zooplankton with respect to population status of river Tapi in North Maharastra region. *J. Curr. Sci*., 5: 749 – 754.
Garcia, J.R., and Lopez, J.M., 1989. Seasonal patterns of phyto plankton productivity, zooplankton
Gayathri, S., Latha, N., Mohan, M.R. 2014.Studies on population dynamics and seasonal abundance of zooplankton community in Doddavoderahallilake,Bangalore. *Int. J. Emerg. Trends Eng.Dev.*,4(1): 50 55.Goswami, A.P., Mankodi, P.C. 2012. Study on zooplankton of freshwater reservoir
George, M. G. (1961): Observation on theRotifers from Shallow Ponds in Delhi. *Curr.*
Giller, P. S. and Malmqvist, Bjorn (2002): The Biology of Streams and Rivers. Oxford University Press Inc. New York. 235pp.

Gopal B., Chauhan M. (2001). South Asian wetlands and their diversity: the role ofmonsoons. - In: Gopal B, Junk WJ,Davis JA (eds) Biodiversity in wetlands: assessment, function and conservation, vol 2., Backhuys Publishers, Leiden, pp257-276.

Gopal B., Zutshi D.P (1998). Fifty years of hydrobiological research in India. – Hydrobiologia 384:267-290.

Goswami A.P (2012). Department of Biology, M.V.M. Science and Home Science College, Rajkot, Gujarat, INDIA.ISCA Journal of Biological Sciences Vol.1 (1), 30-34.

Goswami A.P. and Mankodi P.C., Study on Zooplankton of Fresh Water Reservoir Nyari-II Rajkot district, Gujarat,India, *ISC Journal of Biological Sciences,* 1(1), 30-34(2012)

Guiry, MD, Guiry, GM. *AlgaeBase*, World-wide electronic publication,Galway, National University of Ireland Press, 2011.

Huisman, J., Matthijs, H.C.P. and Visser, P.M(2005). Harmful Cyanobacteria. *Berlin, Springer,* p. 243.

Hujare M S (2008). "Seasonal Variation of Physico-Chemical Parameters in the Perennial Tank of Talsande,Maharashtra" *Ecotoxical. Environ. Monit*., Vol. 18, No. 3,pp. 233-242.16.

Hynes, H.B.N (1971). *The Biology of Polluted Water*. Univ. Toranto Press, Canada pp. 202.

Imoobe, T. O. T. and Adeyinka, M. L. (2010): Zooplankton-based assessment of the trophic state of a tropical forest river. *International Journal of Fisheries and Aquaculture, (I.J.F.A),* **2**(2): 064-070.

Jalilzadeh A.K.K., Yamakanamardi S.M. and AltaffK.,Abundance of zooplankton in three contrasting lake ofMysore city, Karnataka state, India, Sengupta M. andDalwan R (eds.) Proceedings of Taal 2007: The 12 the World lake Conference: 464-469 (2008)

Jana, B.B., 1973. Seasonal periodicity of plankton in a fresh water pond, West Bengal, India, Intl. Rev. Ges. Hydrological. 58: 127 – 143.

Kadam S.S. and Tiwari L.R., Zooplankton Composition in Dahanu Creek-West Coast of India, *Research Journal of Recent Sciences.,* 1(5), 62-65 (2012)

Kaul, V. & A.K. Pandit (1982). Biotic factors and food chain structure in some typical wetlands of Kashmir. *Pollution Research* 1: 49–54.

Kemdirim, E. C. (2000): Diel rhythm of plankton and physico-chemical parameters in Kangimi reservoir, Kaduna State, Nigeria. *Journal of Aquatic Sciences,* **15**: 35-39.

Khan, M.A. (1987). Observations on zooplankton composition, abundance and periodicity in two flood plain lakes of the Kashmir Himalayan valley. *ActaHydrochemicaHydrobiologia*15: 167–174.

Khan, R. A. (2002). The ecology and faunal diversity of two floodplain Ox-bow lakes of south-eastern West Bengal. *Records of the Zoological Survey of India, Occasional Paper No.* 195: 1–57.

Khan, R.A. (2003). Faunal diversity of zooplankton in freshwater wetlands of southeasternWest Bengal. *Records of the Zoological Survey of India, Occasional Paper No.* 204: 1–107.

Koste, W. (1978). *ROTATORIA*. Die RädertiereMitteleuropas, begründet von Max Voigt.ÜberordnungMonogononta. GebrüderBorntraeger, Berlin, Stuttgart. I. Text (673 pp) U. II. Tafelbd. (T. 234).

Kumar, S., and Dutta, S.P.S., 1991. Studies on Phytoplankton population dynamics in Kunjawani pond, Jammu, *J. hydrobiol*., 7(1): 55 – 89.

Lakshmi Ganesan and R.A Khan (2007).Studies on the ecological of zooplankton in a flood plain wetland of WestBengal, India. Proceedings of Taal .The 12th World Lake Conference: 67-73.

Li, Yang; Xu, Linyu and Li, Shun (2009): Water quality analysis of the Songhua River basin using multivariate techniques. *Journal of Water Resources and Protection (JWARP).*

Ludwig, J.A. & J. F. Reynolds (1988). *Statistical Ecology: A Primer on Methods and Computing*. John Wiley & Sons, New York, 337pp.

Magurran, A.E. (1988). *Ecological Diversity and its Measurement*. Croom Helm Limited, London, 179pp.

Mahor R.K., Diversity and seasonal fluctuation ofzooplankton in fresh water reservoir Tighra Gwalior (M.P),*InternationalReseach Journal*, 2(19), 24-25

(2011)22. Krishnamoorthi A. and Selvakumar S., Seasonal fluctuationof zooplankton community in relation to certain physicochemicalparameters of Veeranamlake in Cuddaloredistrict, Tamil Nadu, *Iner. J. Resea. Enviro. Scie. Tech.*,2(2), 22-26 (2012)

Mani, P. and Krishnamoorthy, K., 1989. Ecology of phytoplankton blooms in the velar estuary. East coast of India. *Int. J. Mar. Sci.* 15: 24 – 28.

Manjare S A, Vhanalakar S A and D V Muley (2010). "Water Quality Assesment of Vadgaon tank of Satara(Maharastra), with Special Reference to Zooplankton", *International J. of Advanced Biotechnology and Research,* Vol.1, No. 2, pp. 91-95.

Mann C.J., Wetzel R.G. (1995). Dissolved organic carbon and its utilization in a riverine wetland ecosystem. –Biogeochemistry 31: 99-120.

Marine Biology Organisation (MBO), (2007): ZooplanktonRetrieved Sept. 29. (2006) Retrieved from:http://www.marinebiocom/oceans/zooplankton.asp.62k.

Michael, P. *Ecological Methods for Field and Laboratory Investigations*, TataMcGraw-Hill Publishing Company Ltd., New Delhi, India, 1984.

Michael, R.G. & B.K. Sharma (1988). INDIAN CLADOCERA. (Crustacea: Branchiopoda: Cladocera). *Fauna of India*

Moghraby, A. L. (1977): A study on diapauses of zooplankton in a tropical river – the Blue Nile. *Freshwat. Biol., 7:* 77-117.

Nandan S.N. (1983). Algal flora of polluted waters of Gujarat state Biodiversity Scenario. Discovery Publication House,New Delhi. Pp. 152-170.

National workshop on reservoir fisheries,Special Publication. A.F.S.I. Branc., 3: 103-106.

Naz, S., Musfaqua, N. and Sharminz, (2006):Primary Productivity of an Artificial Lake in

Negi, R. K., Johan, N. S. and Negi, T. (2006):Study of the Physico-chemical Parameters of

Nirmal Kumar – CiniOommen. (2011). Phytoplankton composition in relation to hydrochemical properties of tropicalcommunity wetland. Applied ecology and environmental research 9(3): 279-292.(on line)

Nirmal Kumar J.I., Kumar, R.N., Bhatt, I. (2005). Study of Cultural Eutrophication in relation to the plant diversity ofWetlandRathreshwar in central Gujarat. – In: Aquatic
Okogwu I.O., Seasonal variations of species compositionand abundance of zooplankton in Eboma Lake, FloodplainLake in Nigeria, *Rev. Biol. Trop*., 58(1), 171-182 (2010)20. Chauhan R., Seasonal fluctuation of zooplanktons inRenuka lake Himachala Pradesh, *Utter Pradesh J. Zool*.,113(1), 17-20 (1993)
Oxford and IBH Publishing Co. Pvt. Ltd., NewDelhi. 1158 pp.
Pandey B.N., Jha A.K. and Pandey K., Zooplankton community in relation to certain physico-chemical factors of Kosi, Purnes, Bihar, *Environ. Ecol*., 12, 563-567 (1994)
Pandey, J. and Sharma, M. S. (2003): Env. Sci.Practical and Field Manual, Yash Pub. House,Bikaner.
Parameters of Amarchand Reservoir, SouthernRajasthan. *Poll. Res.,* 13(2): 217-226.

Pathwa D.V. and S. N. Mehotra (1966): Observation on fluctuations in the abundance of plankton in relation to certain hydrobiological condition of river Ganga
Patrick R. P. (1948): Factor affecting the distribution of diatoms Bot. Rev 14(8):473-524.
Pennak R. W. (1962): Quantitative zooplankton sampling in literal vegetation area. Limnol. Oceangr 7:487-489.

Pennake R. W. (1953): freshwater invertebrates of united states 2nd edition John Willey and Sons Inc . 763.
Peumal, P., Swampathkumar, P. andSanthanam, P. K. (1998): Zooplankton ofParangipettal costal water. Monogr. Series,Vol. l. *UGC- SAP, CAS in Marine Biology,*Annamali University, Parangipettai. P. 31.
Philopose M. T. (1960): Freshwater phytoplankton of inland fisheries proc. Sym. Algology ICAR, 272-291.
phytoplankton dynamics and productivityfluctuations in Gandhi Sagar Reservoir.

Pinto A.A.,Zooplankton community structure of two marginal lakes of the river Cuiaba (Mato Grosso, Brazil) with analysis ofRotifera and Cladocera diversity, *Braz. J. Biol*., 63, 1-20 (2003)
Poll. Res., 28 (4): 571-576.
Rajasekar, K.J., Perumal, P., Santhanam, P., 2005. Phytoplankton diversity in the coleroon. Estuary, South East coast of India. *J. Mar. Sci., Biol. Assoc*. India. 47: 127 – 132.
Rajashahi, Bangladesh. *Nat. Env. Poll. Tech*.5(1): 139-141.
Rana B.C., Nirmal Kumar J.I. (1992). Macrophytes and Nutrient study of two wetlands of Guajrat, India. – InternationalJournal Ecology and Environmental Science.18: 195-202.
Ranu (2001): Studies on Toxicity of TextileEffluents to Fresh Water Zooplankton. Ph.D.
Rath, R. K. (2000): Freshwater aquaculture (2nd Ed.) Scientific Publishers (India), Jodhpur. 445.
Rawson, D.S., 1956. Algal indicators of Tropic lake type. Limnology and oceanography. 1: 18 – 25.

Reid, G. K. and Wood, R. D. (1976): Ecology ofInland Waters and Estuaries. *D. Van NostandCompany*, New York.
Research Article Jagadeeshappa K.C et al, Carib.j.SciTech,2013,Vol.1,185-193193
Saha, S.B., Bhattacharya, S.B., and Choudhary, A., 2000. Diversity of phytoplankton of a sewage pollution brackish water tidal ecosystem. *J. Environ Biol.*, 21(1): 9 – 14.
Sanjer, L.R. & U.P. Sharma (1995). Community structure of plankton in Kawar lake wetland, Begusarai, Bihar: II Zooplankton. *Journal of Freshwater Biology* 7: 165–167.
Sarang, N. (2001): Selected Aspects ofLimnology and Productivity of Jaisamand Lake,

Sci., 30: 268-269..Segers, H. (2008). Global diversity of rotifers (Rotifera) in freshwater. *Hydrobiologia*595: 49–59.*and adjacent countries Series*. Published by Zoological Survey of India, Calcutta, 262pp.

Selected Waters of Rajasthan Ph.D. thesisSubmitted to University of Udaipur.
Shaji C. (1990). Studies in algal flora and ecology of some polluted waters in Gujarat Stanley E.H., Ward A.K. (1997).Inorganic nitrogen regimes in an Albama wetland. – Journal of the North American Benthological Society 16:820-832.
Sharma, B. K. & S. Sharma (1999a). Freshwater Rotifers (Rotifera: Eurotatoria). In: *State Fauna Series: Fauna of Meghalaya.* 4(9): 11–161. Zoological Survey of India, Calcutta.
Sharma, B. K. & S. Sharma (1999b). Freshwater Cladocerans (Crustacea: Branchiopoda: Cladocera). In: *State Fauna Series: Fauna of Meghalaya* 4(9): 469 –550. Zoological Survey of India, Calcutta.

Sharma, B. K. & S. Sharma (2009a). Faunal diversity of Cladocera (Crustacea: Branchiopoda) of Loktak Lake (a Ramsar site), Manipur (N.E. India). *Journal of the Bombay Natural History Society* 106(2): 156–161.
Sharma, B. K. (2009a). Diversity of Rotifers (Rotifera: Eurotatoria) of Loktak lake, North-Eastern India. *Tropical Ecology* 50(2): 277–285.
Sharma, B.K. & M. Hussain (2001). Abundance and ecology of Zooplankton in a tropical floodplain lake, Assam (N.E. India). *Ecology, Environment & Conservation* 7(4): 397–403.

Sharma, B.K. & S. Sharma (2000). Freshwater Rotifers (Rotifera: Eurotatoria). In: *State Fauna Series: Fauna of Tripura* 7(4): 163–224. Zoological Survey of India, Calcutta.
Sharma, B.K. & S. Sharma (2001). Biodiversity of Rotifera in some tropical floodplain lakes of the Brahmaputra river basin, Assam (N.E. India). *Hydrobiologia*446/447: 305–313.
Sharma, B.K. & S. Sharma (2005). Faunal diversity of Rotifers (Rotifera: Eurotatoria) of Deeporbeel, Assam (N.E. India) - a Ramsar site. *Journal of the Bombay Natural History Society* 102(2): 169–175.
Sharma, B.K. & S. Sharma (2009b). Diversity of microcrustacea (Crustacea: Branchiopoda) of Loktak lake, a Ramsar site, Manipur, India. *Journal of Threatened Taxa* 1(11): 541–548
Sharma, B.K. (1998). Freshwater Rotifers (Rotifera: Eurotatoria). In: *Fauna of West Bengal. State Fauna Series* 3(11): 341–461. Zoological Survey of India, Calcutta.
Sharma, B.K. (2000a). Rotifers from some tropical flood-plain lakes of Assam (N.E. India). *Tropical Ecology* 41(2): 175–181.
Sharma, B.K. (2000b). Synecology of Rotifers in a tropical floodplain lake of Upper Assam (N. E. India). *The Indian Journal of Animal Sciences* 70: 880–885.
Sharma, B.K. (2005). Rotifer communities of floodplain lakes of the Brahmaputra basin of lower Assam (N.E. India): biodiversity, distribution and ecology. *Hydrobiologia*533: 209–221.

Sharma, B.K. (2009b). Richness, abundance and ecology of Rotifera of Loktak Lake (a Ramsar site), Manipur, Northeastern India. *Ecology, Environment & Conservation* 15(2): 299–306.

Sharma, K. (2007): Recent Limnologicalchanges in Different Parameters of Ranisar
Sharma, L. L. (1980): Some LimnologicalAspects of Udaipur Waters in Comparison to
Sharma, L. L. and Gupta, M. C. (1994):Seasonal Variations in Selected Limnochemical
Sharma, L. L., Sharma, M. S., Lodha, R. M. andDurve, V. S. (1996): J.S.S. Report on Present

Sharma, M. S., Sharma, L. L. and Durve, V. S.(1984): Eutrophication of Lake Pichhola in
Sharma, R., Sharma V., Malera, H. andSharma, M. S. (2009): Study of Microbiology of
Sharma, V. (2007): Biodiversity of Planktonicand Littoral Cladocerans in Water Bodies of
Sharma, V., Ridhhi, S., Malera, H. and Sharma,M. S. (2009): Zooplanktonic Diversity and
Shekhawat, P. S. (1997): Some Aspects ofHydrobiology of a Temple Tank of Udaipur forAssessing it's Trophic status for Fish Rearing.M.Sc. Thesis Submitted to RAU, Bikaner.
Singh, S. P., Pathak, D. and Singh, R., (2002) Eco. Environ. Cons 8: 289.

Sinha B. and Islam M.R., Seasonal variation in zooplankton population of two lentic bodies and Assam State Zoo cumBotanical garden, Guwahati, Assam, *Eco. Environ. Cons*.,8, 273-278 (2002)
Solomon, I. J. J. (1994): Current Trends inEutrophication of the Lakes PichholaandFatehsagar. M.Sc. Thesis Submitted to RAU,Bikaner.
Soruba R., Ecology of water bodies formed by limestonemining in and around Ariyalure, Ph.D., Thesis,Bharathidasan University, (2002)
South Rajasthan. Ph.D. Thesis Submitted toM.L. Sukhadia University, Udaipur.
Status of Udaipur lakes. 1-116pp.
Stronkhorst J., Brils J., Batty J., Coquery M., Gardner M., Mannio J., O'Donnell C., SteenwijkJ.andFrintrop P. (2004).Discussion document on Sediment Monitoring Guidance for the EU.
studies, Environm. Pub, Karad, India, 1984.
Sukunan, V.V., 1980. Seasonal fluctuations of plankton of NagarajunaSagar reservoir, Andhra Pradesh, India. *J. Inland Fish*. Soc. India. 12 : 79 – 91.
Sumitra, M., Sharma, R., Sharma V. andSharma, M. S. (2007). Trophic Status of LakePichhola in Relation to Physico-chemicalCharacteristics of its Water. *Nationalsymposium on Limnology*. 244-248pp.
Talwar, P.K., Jhingran, A .G. (1991): Inland Fishesof India and adjacent countries Vol. 1 and 2.
thesis Submitted to MLSU, Udaipur.26. Rao, K. S. and Choubey, U. (1990): Studies on
Tiwari, A., and Chauthan, S.V.S., 2006. Seasonal phytoplankton diversity of Kitham Lake. *Agra J. Environ. Biol*. 27: 35 – 38.
Trivedy, RK, Goel, PK. Chemical and biological methods for water pollution

Trophic Status of Mahi Dam in Relation toPhysico-chemical Characteristics of its Water.
Tsai C., Ali Y. (1997). Openwater fisheries of Bangladesh. – The University Press Limited, Dhaka
Udaipur (Rajasthan). M.Sc. Thesis Submittedto MPUAT, Udaipur (Rajasthan).
Udaipur Lakes in Relation to DifferentLimnological Parameter. *Poll. Res.,* 28(4): 565-569.
Udaipur, Rajasthan. *Poll. Res.,* 3(2): 39-44.
Upadhayay, B. L. and Dwivedi, D. D. (2006): Impact of Human Activities on PhysicochemicalCharacterstics of Mansi Ganga wateratGoverdhan (Mathura). *Him.J.Env.Zool.*,20(1):131-133.

Verma M C, Singh S and Thakur P (2001).“Ecology of a Perennial Wetland. An Overview of Limnobiotic Status”, *J.Environ.Poll.,* Vol. 8(1): 53-59.
Vijaylaxmi, P. (2003): Biological Productivity ofTwo dry Bundhs of Udaipur, Rajasthan inRelation to Growth and Diet selectivity ofMajor Carp Fry and Fingerlings. Ph.D. ThesisSubmitted to M.L. SukhadiaUniversity,Udaipur.
Vyas, L. N. (1968): Studies on thePhytoplankton Ecology of PichholaLakeUdaipur. Proc. *Symp. Recent. Adv. Tro.Ecology.,* 334-337.
W.H.O. (1973): Guidelnes for Drinking WaterQuality, World Health Organisation, Geneva
Water of Pongdam Reservoir, HimachalPradesh: A Ramsar site. *Him.I. Env. Zool*., 20(2): 247-256

Welch P (1952). “Limnologicalmethods.TheBlankston Co., Philadelphia. Toronta, Wuhrmann, K 1938, Lake”,Eutrophication and its Control. In Proc. Shiga Conference on Conservation and Management of World Environment:26-37.
Wetlands Ecol Manage 11: 273-280.
Whitton, B.A. and Patts, M. (2000). The Ecology of Cyanobacteria. Kluwer Academic Publishers, Netherlands.
Zafaar, A.R., 1986, Seasonality in phytoplankton in some South Indian lakes Hydrobiol. 138: 117 – 187.